

On the Stability of Planetary Circular Orbits in Noncommutative Spaces

Kourosh Nozari and Siamak Akhshabi

*Department of Physics, Faculty of Basic Science,
University of Mazandaran,
P. O. Box 47416-1467, Babolsar, IRAN
e-mail: knozari@umz.ac.ir*

Abstract

We investigate the effects of space noncommutativity and the generalized uncertainty principle on the stability of circular orbits of particles in both a central force potential and Schwarzschild spacetime. We find noncommutative form of the effective potential which up to first order of noncommutativity parameter contains an angular momentum dependent extra term. This angular momentum dependent extra term affects the stability of circular orbits in such a way that the radius of a stable circular orbit in noncommutative space is larger than its commutative counterpart. In the case of large angular momentum, the condition for stability of circular orbits in noncommutative space differs considerably from commutative case.

PACS: 02.40.Gh, 04.60.-m, 03.65.Sq, 91.10.Sp

Key Words: Noncommutative Geometry, Generalized Uncertainty Principle, Planetary Orbits

1 Introduction

An important consequence of quantum gravity scenarios such as string theory is the possible noncommutativity of spacetime structure at short distances[1-7]. This noncommutativity leads to the modification of Heisenberg uncertainty relations in such a way that prevents one from measuring positions to better accuracies than the Planck length. In low energy limit, these quantum gravity effects can be neglected, but in circumstances such as very early universe or in the strong gravitational field of a black hole one has to consider these effects. The modifications induced by the generalized uncertainty principle on the classical orbits of particles in a central force potential firstly has been considered by Benczik *et al*[8]. The same problem has been considered within noncommutative geometry by Mirza and Dehghani[9]. The main consequence of these two investigation is the constraint imposed on the minimal observable length and noncommutativity parameter in comparison with observational data of Mercury. Here we are going to proceed one more step in this direction. We study the effects of the generalized uncertainty principle and space noncommutativity on the stability of circular orbits of particles in a central force potential and also Schwarzschild geometry. We obtain a noncommutative effective potential in each case, where up to first order of noncommutativity parameter, there is an extra angular momentum dependent term which affects the conditions for stability of circular orbits. For large values of angular momentum, the effect of space noncommutativity is considerable.

2 Preliminaries

In usual quantum mechanics one has the following canonical commutation relations:

$$[\tilde{x}_i, \tilde{x}_j] = 0, \quad [\tilde{p}_i, \tilde{p}_j] = 0, \quad [\tilde{x}_i, \tilde{p}_j] = i\hbar\delta_{ij} \quad (1)$$

From a string theory point of view, existence of a minimal length scale can be addressed in the following generalized uncertainty principle[10,11]

$$\delta\tilde{x}_i\delta\tilde{p}_i \geq \frac{\hbar}{2}\left(1 + \beta(\delta\tilde{p}_i)^2 + \gamma\right), \quad (2)$$

where β is string theory parameter related to minimal length[11]. Since we are going to deal with absolutely minimum position uncertainty, we set $\gamma = \beta\langle\tilde{p}\rangle^2$ and therefore the corresponding canonical commutation relation becomes

$$[\tilde{x}_i, \tilde{p}_j] = i\hbar\delta_{ij}(1 + \beta\tilde{p}^2), \quad (3)$$

where $\tilde{p}^2$ is the squared value of momentum. Now, canonical commutation relations in commutative space with GUP becomes

$$[\tilde{x}_i, \tilde{x}_j] = 0, \quad [\tilde{x}_i, \tilde{p}_j] = i\hbar\delta_{ij}(1 + \beta\tilde{p}^2), \quad [\tilde{p}_i, \tilde{p}_j] = 0. \quad (4)$$

In a noncommutative space, the set of relations (1) should be changed to the following relations

$$[\tilde{x}_i, \tilde{x}_j] = i\theta_{ij}, \quad [\tilde{p}_i, \tilde{p}_j] = 0, \quad [\tilde{x}_i, \tilde{p}_j] = i\hbar\delta_{ij} \quad (5)$$

where θ_{ij} is an anti-symmetric matrix which determines the fundamental cell discretization of space-time in the same way as the Planck constant $\hbar$ discretizes the phase space. The elements of θ_{ij} have the dimension of $(length)^2$. In the classical limit, the commutators should be replaced by corresponding Poisson brackets

$$\frac{1}{i\hbar}[\tilde{A}, \tilde{B}] \longrightarrow \{\tilde{A}, \tilde{B}\} \quad (6)$$

and therefore in this limit the commutation relations become as follows

$$\{\tilde{x}_i, \tilde{x}_j\} = \alpha_{ij}, \quad \{\tilde{p}_i, \tilde{p}_j\} = 0, \quad \{\tilde{x}_i, \tilde{p}_j\} = \delta_{ij} \quad (7)$$

here $\alpha_{ij} = \frac{\Theta_{ij}}{\hbar}$. The Poisson bracket should possess the same properties as the quantum mechanical commutator, namely, it should be bilinear, anti-symmetric and should satisfy the Leibniz rules and the Jacobi identity. The general form of the Poisson brackets for this deformed version of classical mechanics can be written as follows [8,9]:

$$\{A, B\} = \left[\frac{\partial A}{\partial \tilde{x}_i} \frac{\partial B}{\partial \tilde{p}_j} - \frac{\partial A}{\partial \tilde{p}_i} \frac{\partial B}{\partial \tilde{x}_j} \right] \{\tilde{x}_i, \tilde{p}_j\} + \frac{\partial A}{\partial \tilde{x}_i} \frac{\partial B}{\partial \tilde{x}_j} \{\tilde{x}_i, \tilde{x}_j\} \quad (8)$$

where repeated indices are summed. There is a new coordinate system which can be defined with the following transformations

$$x_i = \tilde{x}_i + \frac{1}{2}\alpha_{ij}\tilde{p}_j, \quad p_i = \tilde{p}_i \quad (9)$$

where the new variables satisfy the usual canonical brackets

$$\{x_i, x_j\} = 0, \quad \{p_i, p_k\} = 0, \quad \{x_i, p_j\} = \delta_{ij}. \quad (10)$$

With these preliminaries, we have all prerequisites to investigate particle orbits and their stability within noncommutative geometry and the generalized uncertainty principle.

3 Central Force Potential

In noncommutative space, for a particle in a central force potential the Hamiltonian has the following form

$$H = \frac{\tilde{p}^2}{2m} + V(\tilde{r}), \quad \tilde{r} = \sqrt{\tilde{x}_i \tilde{x}_j} \quad (11)$$

Using coordinates transformation given as (9), we find

$$\begin{aligned} V(\tilde{r}) &= V\left(\sqrt{(x_i - \frac{\alpha_{ij} p_j}{2})(x_i - \frac{\alpha_{ik} p_k}{2})}\right) \\ &= V(r) + \frac{\vec{\alpha} \times \vec{p}}{2} \cdot \vec{\nabla} V(r) + O(\alpha^2) \\ &= V(r) - \frac{(\vec{\alpha} \cdot \vec{L})}{2r} \frac{\partial V}{\partial r} + O(\alpha^2), \end{aligned} \quad (12)$$

where $\alpha_{ij} = \epsilon_{ijk} \alpha_k$ and $\vec{L} = \vec{r} \times \vec{p}$. In which follows, we first determine the effects of space noncommutativity on the planetary orbits and then we investigate the stability of circular orbits in a general central force potential.

3.1 Particle Orbits

The effect of space noncommutativity on orbital motions of particles in a central force potential has been investigated by Mirza and Dehghani[9]. For an inverse squared force(Coulomb force) as $F = -\frac{k}{r^2}$, the Hamiltonian up to the first order of α becomes

$$H = \frac{p^2}{2m} - \frac{k}{r} - \frac{k}{r^3} \left(\frac{\vec{\alpha} \cdot \vec{L}}{2} \right). \quad (13)$$

The new term in the Hamiltonian is small and its effects can be obtained by standard perturbation theory, however it causes a time dependent angular momentum. We assume that the time variation of the $\vec{L}$ is so small that in a short time interval (for example one century for Mercury) it could be taken as a constant of the motion. In this way, one can put a bound on the value of α by comparing the results of the perturbing term with the experimental value of the precession of the perihelion of Mercury[8,9]. For the Kepler problem it is well known that the bounded orbits are closed, which is a result of the following integral,

$$\Delta\varphi^{(0)} = -2 \frac{\partial}{\partial L} \int_{r_{min}}^{r_{max}} \sqrt{2m[E - V(r)] - \frac{L^2}{r^2}} dr = 2\pi, \quad (14)$$

where

$$E = \frac{1}{2}m(\dot{r}^2 + r^2\dot{\varphi}^2) + V(r), \quad L = mr^2\dot{\varphi}.$$

By perturbing the potential with a small term $V \rightarrow V + \delta V$ and expanding the integral up to the first order in δV , we find

$$\Delta\varphi = \Delta\varphi^{(0)} + \Delta\varphi^{(1)} \quad (15)$$

where

$$\Delta\varphi^{(1)} = \frac{\partial}{\partial L} \int_0^\pi \left(\frac{2mr^2}{L} \right) \delta V d\varphi. \quad (16)$$

After a straightforward calculation, we arrive at the following relation

$$\Delta\varphi^{(1)} = \left(\frac{2\pi k^2 m^2 \cos(\psi)}{L^3} \right) \alpha \quad (17)$$

where ψ is the angle between $\vec{\alpha}$ and $\vec{L}$. This is the effect of space noncommutativity on the orbits of particles in a central force potential. Comparison with observational data of Mercury sets a bound on the noncommutativity parameter[9].

3.2 Stability of the Circular Orbits

Now we consider a central force with the following form

$$F = -\frac{k}{r^n}. \quad (18)$$

In this case, the effective potential up to first order of α can be written as

$$V_{eff}(\tilde{r}) = -\frac{k}{n-1} \frac{1}{r^{n-1}} + \frac{L^2}{2mr^2} - \frac{k}{r^{n+1}} \left(\frac{\vec{\alpha} \cdot \vec{L}}{2} \right). \quad (19)$$

Here m is the mass of the particle and $L = mr^2\dot{\varphi}$. The particle could have a stable circular orbit at $r = \rho$ if the effective potential has a real minimum at ρ . In other words, the effective potential should satisfy the following conditions to have stable circular orbit at ρ

$$\frac{\partial V_{eff}}{\partial r} \Big|_{r=\rho} = 0, \quad \text{and} \quad \frac{\partial^2 V_{eff}}{\partial r^2} \Big|_{r=\rho} > 0. \quad (20)$$

With effective potential as (19), we find

$$\frac{\partial V_{eff}}{\partial r} \Big|_{r=\rho} = \frac{k}{\rho^n} - \frac{L^2}{m\rho^3} + \left(\frac{\vec{\alpha} \cdot \vec{L}}{2} \right) \frac{k(n+1)}{\rho^{n+2}} = 0, \quad (21)$$

and

$$\frac{\partial^2 V_{eff}}{\partial r^2} \Big|_{r=\rho} = -\frac{nk}{\rho^{n+1}} + \frac{3L^2}{m\rho^4} - \left(\frac{\vec{\alpha} \cdot \vec{L}}{2} \right) \frac{k(n+1)(n+2)}{\rho^{n+3}} > 0. \quad (22)$$

These two relations can be rewritten as follows respectively

$$k\rho^2 - \frac{L^2}{m}\rho^{n-1} + \left(\frac{\vec{\alpha} \cdot \vec{L}}{2}\right)k(n+1) = 0, \quad (23)$$

and

$$-nk\rho^2 + \frac{3L^2}{m}\rho^{n-1} - \left(\frac{\vec{\alpha} \cdot \vec{L}}{2}\right)k(n+1)(n+2) > 0. \quad (24)$$

From equation (23) we find

$$\frac{L^2}{m}\rho^{n-1} = k\rho^2 + \left(\frac{\vec{\alpha} \cdot \vec{L}}{2}\right)k(n+1). \quad (25)$$

Using this result in (24), we obtain the following condition for stability of circular orbits in central force of the type given as (18)

$$(3-n)k\rho^2 - \left(\frac{\vec{\alpha} \cdot \vec{L}}{2}\right)k(n+1)(n-1) > 0, \quad (26)$$

or

$$(3-n)k\rho^2 - \frac{k(n+1)(n-1)L\alpha \cos(\psi)}{2} > 0, \quad (27)$$

where ψ is the angle between $\vec{\alpha}$ and $\vec{L}$. In the absence of space noncommutativity, i.e., when $\alpha = 0$, we find the usual result of $n < 3$, as it is well known from classical mechanics. In the presence of space noncommutativity, the third term in effective potential (19) causes a time dependent angular momentum which affects the stability of circular orbits. However the time variation of $\vec{L}$ is so small that in a short time interval it could be taken as a constant of the motion. Figure 1 shows the conditions for stability of circular orbits in a central force of the type given in (18). As this figures shows, for small values of L , the condition for stability of circular orbits is the same as classical commutative result, i. e. , $n < 3$. When L increases, the condition for stability differs considerably relative to commutative case. In this case n can attains non-integer values. This feature may be interpreted as the possible deviation from inverse square law. In another words, possible deviation from inverse square law can be interpreted as a consequence of space noncommutativity.

3.3 Stability of Circular Orbits in Yukawa Potential

As another specific example, here we consider the Yukawa potential and investigate the stability of its circular orbits in noncommutative space. Yukawa potential is defined as follows

$$V(r) = -\frac{k}{r}e^{-r/a} \quad k > 0, \quad a > 0. \quad (28)$$

Using equation (12), the corresponding potential in noncommutative space becomes

$$V(\tilde{r}) = -\frac{k}{r}e^{-r/a} - k\left(\frac{\vec{\alpha} \cdot \vec{L}}{2}\right)\left(\frac{1}{ar^2} + \frac{1}{r^3}\right)e^{-r/a}, \quad (29)$$

and therefore the effective potential is given by

$$V_{eff}(\tilde{r}) = -\frac{k}{r}e^{-r/a} - k\left(\frac{\vec{\alpha} \cdot \vec{L}}{2}\right)\left(\frac{1}{ar^2} + \frac{1}{r^3}\right)e^{-r/a} + \frac{L^2}{2mr^2}. \quad (30)$$

By applying the conditions (20) to this $V_{eff}(\tilde{r})$, a straightforward calculation gives the following condition for stability of circular orbits in Yukawa potential

$$3 + \rho \frac{F'(\rho)}{F(\rho)} > 0, \quad (31)$$

where $F(r) = -\frac{\partial V}{\partial r}$ and $F'(r) = \frac{\partial F}{\partial r}$. A simple calculation gives

$$3 - \frac{\left[\frac{2}{\rho^2} + \frac{2}{a\rho} + \frac{1}{\rho^2}\right] + \left[\frac{\vec{\alpha} \cdot \vec{L}}{2}\right]\left[\frac{12}{\rho^4} + \frac{12}{a\rho^3} + \frac{5}{a^2\rho^2} + \frac{1}{a^3\rho}\right]}{\left[\frac{1}{a\rho} + \frac{1}{\rho^2}\right] + \left[\frac{\vec{\alpha} \cdot \vec{L}}{2}\right]\left[\frac{3}{\rho^4} + \frac{3}{a\rho^3} + \frac{1}{a^2\rho^2}\right]} > 0, \quad (32)$$

or

$$(a^2\rho^3 + a^3\rho^2 - a\rho^4) - \left(\frac{\vec{\alpha} \cdot \vec{L}}{2}\right)(\rho^3 + 2a\rho^2 + 3a^2\rho + 3a^3) > 0. \quad (33)$$

This is the condition for stability of circular orbits of Yukawa potential in noncommutative space. If $\alpha = 0$, this relation reduces to the following condition

$$a\rho + a^2 - \rho^2 = 0, \quad (34)$$

which leads to the condition

$$\rho \leq 1.62a$$

for stability of circular orbits. This result is well known from classical mechanics.¹

¹A digression to a newly proposed scenario may be helpful here. Equation (34) has the following solutions

$$\rho = \frac{a}{2} \pm \sqrt{\frac{a^2}{4} + 1} = a\frac{1}{2}\left[1 \pm \sqrt{\frac{5}{4}}\right]$$

where in the language of E-infinity theory[12,13] can be interpreted as follows

$$\rho_+ = a\frac{1}{2}[1 + \sqrt{5}] = a\left(\frac{1}{\phi}\right)$$

and

$$\rho_- = a\frac{1}{2}[1 - \sqrt{5}] = -a(\phi)$$

where ϕ is the golden mean. Note also that in this framework

$$\rho \leq (1.618033969)a.$$

Figure 2 shows the condition for stability of circular orbits of particles in Yukawa potential. As this figure shows, for small values of L , the condition for stability is essentially the same as commutative case. For large L however the situation is different from commutative case. For large L , we see that the condition can change to, for example,

$$\rho \leq 1.414213562a.$$

Note that since noncommutativity parameter is very small, the role played by $\vec{L}$ is enhanced. But we know that presence of $\vec{L}$ is due to non-vanishing value of α , that is, space noncommutativity. Thus, we can conclude that space noncommutativity (up to first order of noncommutativity parameter) causes $\vec{L}$ -dependent terms in orbital equations which these terms are important only in large values of L .

4 Stability of Circular Orbits in Schwarzschild Geometry

In this section we investigate the existence and stability of circular orbits of particles in noncommutative Schwarzschild geometry. We first obtain commutative effective potential and then generalize it to noncommutative case.

Consider the geometry of Schwarzschild spacetime with the following metric(with $c = 1$)

$$ds^2 = -(1 - \frac{2GM}{r})dt^2 + (1 - \frac{2GM}{r})^{-1}dr^2 + r^2d\Omega^2 \quad (35)$$

The geodesics of this spacetime are given by the following equations[14]

$$\frac{d^2t}{d\lambda^2} + \frac{2GM}{r(r-2GM)} \frac{dr}{d\lambda} \frac{dt}{d\lambda} = 0 \quad (36)$$

$$\begin{aligned} \frac{d^2r}{d\lambda^2} + \frac{GM}{r^3}(r-2GM)(\frac{dt}{d\lambda})^2 - \frac{GM}{r(r-2GM)}(\frac{dr}{d\lambda})^2 \\ - (r-2GM)\left[(\frac{d\vartheta}{d\lambda})^2 + \sin^2(\vartheta)(\frac{d\varphi}{d\lambda})^2\right] = 0 \end{aligned} \quad (37)$$

$$\frac{d^2\vartheta}{d\lambda^2} + \frac{2}{r} \frac{d\vartheta}{d\lambda} \frac{dr}{d\lambda} - \sin(\vartheta) \cos(\vartheta) (\frac{d\varphi}{d\lambda})^2 = 0 \quad (38)$$

$$\frac{d^2\varphi}{d\lambda^2} + \frac{2}{r} \frac{d\varphi}{d\lambda} \frac{dr}{d\lambda} + 2 \cot(\vartheta) \frac{d\vartheta}{d\lambda} \frac{d\varphi}{d\lambda} = 0 \quad (39)$$

where λ is an affine parameter. There are four conserved quantities associated with Killing vectors: magnitude of angular momentum(one component), direction of angular momentum(two components) and energy. The two Killing vectors which lead to conservation

of the direction of angular momentum imply that $\vartheta = \frac{\pi}{2}$. The other two Killing vectors are corresponding to energy and the magnitude of the angular momentum. The Killing vector associated with energy is ∂_t or

$$K_\mu = \left(-\left(1 - \frac{2GM}{r}\right), 0, 0, 0 \right) \quad (40)$$

and for the magnitude of angular momentum the Killing vector is ∂_φ or

$$L_\mu = \left(0, 0, 0, r^2 \sin^2(\vartheta) \right). \quad (41)$$

So, along the geodesics the two corresponding conserved quantities are

$$\left(1 - \frac{2GM}{r}\right) \frac{dt}{d\lambda} = E \quad (42)$$

and

$$r^2 \frac{d\varphi}{d\lambda} = L \quad (43)$$

respectively, where E and L are energy and angular momentum of the particle per its unit mass.

Along an affinely parameterized geodesic (timelike, spacelike or null), the scalar quantity

$$\varepsilon = -u^\alpha u_\alpha$$

is a constant since

$$\frac{d\varepsilon}{d\lambda} = (u^\alpha u_\alpha)_{;\beta} u^\beta = (u_{;\beta}^\alpha u^\beta) u_\alpha + u^\alpha (u_{\alpha;\beta} u^\beta) = 0.$$

If we chose λ to be proper time or proper distance, then we find $\varepsilon = \pm 1$. For a null geodesic $\varepsilon = 0$. Generally, one can write

$$\varepsilon = -g_{\mu\nu} \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda} \quad (44)$$

We chose $\lambda = \tau$ (the proper time) and expand this equation. In equatorial plane ($\vartheta = \frac{\pi}{2}$) we find

$$-\left(1 - \frac{2GM}{r}\right) \left(\frac{dt}{d\lambda}\right)^2 + \left(1 - \frac{2GM}{r}\right)^{-1} \left(\frac{dr}{d\lambda}\right)^2 + r^2 \left(\frac{d\varphi}{d\lambda}\right)^2 = -\varepsilon \quad (45)$$

where multiplying by $\left(1 - \frac{2GM}{r}\right)$ and using equations (48) and (49), we obtain

$$-E^2 + \left(\frac{dr}{d\lambda}\right)^2 + \left(1 - \frac{2GM}{r}\right) \left(\frac{L^2}{r^2} + \varepsilon\right) = 0 \quad (46)$$

This relation can be rewritten as follows

$$\frac{1}{2} \left(\frac{dr}{d\lambda}\right)^2 + V_{eff}(r) = \frac{1}{2} E^2 \quad (47)$$

where we have defined

$$V_{eff}(r) = \frac{1}{2}\varepsilon - \varepsilon \frac{GM}{r} + \frac{L^2}{2r^2} - \frac{GML^2}{r^3} \quad (48)$$

which is the *effective potential* in commutative Schwarzschild spacetime. Figure 3 shows the variation of this effective potential versus radius and for different angular momentum. In terms of the Schwarzschild radius, one can rewrite this equation as follows

$$V_{eff}(r) = \frac{1}{2}\varepsilon - \varepsilon \frac{GM}{r} + \frac{L^2}{2r^2} \left(1 - \frac{r_s}{r}\right) \quad (49)$$

where r_s is the Schwarzschild radius. We use this relation in our forthcoming arguments.

In classical general relativity, the particle could have a circular orbit at r_c if

$$\left(\frac{dV}{dr}\right)_{r=r_c} = 0$$

which leads to the following condition

$$\varepsilon GM r_c^2 - L^2 r_c + 3GML^2 \gamma = 0 \quad (50)$$

where $\gamma = 0$ in Newtonian regime and $\gamma = 1$ in general relativity[14]. Let us consider two possible cases. Firstly, for $\gamma = 1$ and $\varepsilon = 0$ (photons) we have

$$r_c = 3GM \quad (51)$$

Secondly, for $\gamma = 1$ and $\varepsilon = 1$ (massive particles) we find

$$r_c = \frac{L^2 \pm \sqrt{L^4 - 12G^2 M^2 L^2}}{2GM} \quad (52)$$

For $L \rightarrow \infty$, there exist a stable circular orbit at $\frac{L^2}{GM}$ which goes farther and farther away and an unstable one at $3GM$. For small L , at $L = \sqrt{12}GM$ two circular orbits coincide at $r_c = 6GM$ and disappear entirely for smaller L . Therefore, $6GM$ is the smallest possible radius of a stable circular orbit in Schwarzschild metric. In brief, Schwarzschild solution in commutative space possesses stable circular orbits for $r \geq 6GM$ and unstable ones for $3GM < r < 6GM$. Now we consider the effects of space noncommutativity on the stability of circular orbits in Schwarzschild geometry.

5 Noncommutative Space Considerations

In Schwarzschild spacetime, with $\varepsilon = 1$ (the case of timelike geodesics) we find from (48)

$$V(r) = \frac{1}{2} - \frac{GM}{r} + \frac{L^2}{2r^2} - \frac{GML^2}{r^3}. \quad (53)$$

Considering the effects of space noncommutativity as given by (12), this relation changes to the following form

$$V(\tilde{r}) = \frac{1}{2} - \frac{GM}{r} + \frac{L^2}{2r^2} - \frac{GML^2}{r^3} - \frac{L\theta \cos \psi}{2} \left[\frac{GM}{r^3} - \frac{L^2}{r^4} + \frac{3GML^2}{r^5} \right] \quad (54)$$

where ψ is the angle between $\vec{L}$ and $\vec{\theta}$. This noncommutative effective potential has been plotted in figures 4 and 5 for two different values of angular momentum. As these figures show, increasing the value of the particle angular momentum increases the difference between commutative and noncommutative effective potentials. In another words, large angular momentum enhances the effect of space noncommutativity. So space noncommutativity couples with angular momentum. This point has its origin on the algebraic structure of theory on noncommutativity level.

The particle could have a circular orbit at r_c if

$$\left(\frac{dV}{dr} \right)_{r=r_c} = 0$$

or

$$\frac{GM}{r_c^2} - \frac{L^2}{r_c^3} + \frac{3GML^2}{r_c^4} - \frac{L\theta \cos \psi}{2} \left[\frac{-3GM}{r_c^4} + \frac{4L^2}{r_c^5} - \frac{15GML^2}{r_c^6} \right] = 0 \quad (55)$$

This can be simplified to find

$$GMr_c^4 - L^2r_c^3 + 3GML^2r_c^2 - \frac{L\theta \cos \psi}{2} \left[-3GMr_c^2 + 4L^2r_c - 15GML^2 \right] = 0. \quad (56)$$

This is the condition for existence of circular orbits in noncommutative Schwarzschild geometry up to first order of noncommutativity parameter. Comparing this equation with equation (50)(with $\gamma = 1$ and $\epsilon = +1$) for commutative case, shows the importance of noncommutativity effect. Since noncommutativity parameter is very small, this effect can be neglected in ordinary circumstances but for large angular momentum the situation differs considerably. The dependence of this modification to angular momentum is a pure noncommutative effects which goes back to the algebraic structure of the theory.

The condition for the stability of the circular orbits is

$$\left(\frac{\partial^2 V}{\partial r^2} \right)_{r=r_c} > 0 \quad (57)$$

Applying this condition to the potential (54), we find

$$\frac{-2GM}{r_c^3} + \frac{3L^2}{r_c^4} - \frac{12GML^2}{r_c^5} - \frac{L\theta \cos \psi}{2} \left[\frac{12GM}{r_c^5} - \frac{20L^2}{r_c^6} + \frac{90GML^2}{r_c^7} \right] > 0 \quad (58)$$

Combining this equation with equation (55), we find the following condition for the stability of the circular orbits in noncommutative space

$$GMr_c^4 - 3GML^2r_c^2 - \frac{L\theta \cos(\psi)}{2} [3GMr_c^2 - 8L^2r_c + 45GML^2] > 0 \quad (59)$$

Figure 6 shows the condition for stability of circular orbits of particles in Schwarzschild geometry and in the presence of space noncommutativity. In this figure we have plotted the left hand side of relation (58) versus $r = GM$. As this figure shows, in the case of noncommutative space, particles could have stable circular orbits for $r \geq (6.139234690)GM$ which is greater than commutative result of $r \geq 6GM$. Therefore, stable circular orbits have greater radius in noncommutative spaces.

6 The effect of GUP

Now we consider the effect of the generalized uncertainty principle on the stability of circular orbits of particles in Schwarzschild spacetime. With the generalized uncertainty principle, the standard commutation relations transform to the following general form [6,12]

$$[x_i, p_j] = i\hbar(\delta_{ij} + \beta p^2 \delta_{ij} + \beta' p_i p_j), \quad (60)$$

$$[p_i, p_j] = 0, \quad (61)$$

and

$$[x_i, x_j] = i\hbar \frac{(2\beta - \beta') + (2\beta + \beta')\beta p^2}{(1 + \beta p^2)} (p_i x_j - p_j x_i). \quad (62)$$

we set $\beta' = 0$ so the corresponding Poisson brackets are

$$\{x_i, p_j\} = \delta_{ij}(1 + \beta p^2), \quad \{p_i, p_j\} = 0, \quad \{x_i, x_j\} = 2\beta(p_i x_j - p_j x_i) \quad (63)$$

The modified canonical equations now take the following forms

$$\dot{x}_i = \{x_i, H\}, \quad \dot{p}_i = \{p_i, H\} \quad (64)$$

where

$$H = \frac{p^2}{2m} + V(r) \quad (65)$$

is the Hamiltonian of the system. The deformed angular momentum which is given by

$$L_{ij} = \frac{x_i p_j - x_j p_i}{(1 + \beta p^2)} \quad (66)$$

is conserved due to the rotational symmetry of the Hamiltonian. For the circular orbits of particles in a general central force problem we have the constraints $\dot{r} = 0$ and $\dot{p}_r = 0$ [8]. These conditions will lead us to

$$L = \frac{pr}{(1 + \beta p^2)}. \quad (67)$$

The radius of orbit r and the magnitude of the momentum $p = \sqrt{p^2}$ are now the constants of the motion .

Now in Schwarzschild spacetime the effective potential is given by equation (53). We substitute the value of L from (67) in this potential to find

$$V(\tilde{r}) = \frac{1}{2} - \frac{GM}{r} + \frac{p^2}{2(1 + \beta p^2)} - \frac{GMp^2}{r(1 + \beta p^2)} \quad (68)$$

The radius of the circular orbits will be given by

$$r_c = 1/2 \frac{GM(2 + 4\beta p^2 + 2\beta^2 p^4 + 3p^2)}{p^2} \quad (69)$$

so the condition for a circular orbit with radius r to be stable in the framework of GUP is

$$-2GMr_c^2 + \frac{3p^2 r_c^3}{(1 + \beta p^2)^2} - \frac{12GMp^2 r_c^3}{(1 + \beta p^2)^2} \geq 0 \quad (70)$$

We emphasize that presence of β is a quantum gravitational effect which has origin on the fractal structure of spacetime at very short distances(string scale). The situation is much similar to the case presented in figure 6 with dotted curve. The effect of space noncommutativity is much similar to the effect of generalized uncertainty principle. These two concept are common features of quantum gravity era where there exist a minimal observable length of the order of Planck length.

7 Summary and Conclusion

In this paper we have studied the effects of space noncommutativity and the generalized uncertainty principle on the stability of circular orbits of particles in both a central force potential and Schwarzschild geometry. In the case of central force potential, we have shown that when the angular momentum of the particle is small, the effect of space noncommutativity can be neglected. In this case for a central force of the type $F(r) = -\frac{k}{r^n}$, we find $n < 3$ which is condition for stability of circular orbits in commutative classical mechanics. The situation however differs considerably with classical prescription in the case of large angular momentum. In this case, the condition for stability of circular

orbits differs considerably with its classical commutative counterpart. Since the effect of space noncommutativity in first order approximation is the presence of angular momentum in orbital equations, we conclude that space noncommutativity changes conditions for stability of the circular orbits in an angular momentum dependent manner. For clarifying our findings, we have considered the case of Yukawa potential as an example. For the case $n = 2$ in $F(r) = -\frac{k}{r^n}$, from a more deeper point of view, all of our calculations show the implicit departure from inverse squared law due to space noncommutativity. This feature provides an operative approach for investigation of the departure from inverse square law in electricity and also planetary physics.

we have studied the effect of space noncommutativity and the generalized uncertainty principle on the stability of circular orbits of particles in Schwarzschild geometry. We have found the effective potential in the case of noncommutative Schwarzschild space. Again, when the angular momentum of the particle is small, the effect of space noncommutativity can be neglected. In this case we find $r \geq 6GM$ for the stability of circular orbits in commutative Schwarzschild space. The situation differs considerably with commutative prescription in the case of large angular momentum. In this case the condition for stability of circular orbits for $L = 5$ (in arbitrary units) becomes $r \geq (6.139234690)GM$ which differs from commutative result, $r \geq 6GM$. Since the effect of space noncommutativity in first order approximation is the presence of angular momentum in orbital equations, we conclude that space noncommutativity changes conditions for stability of the circular orbits in a angular momentum dependent manner. This feature provides an operative approach for investigation of the spacetime geometry, although this effects are so small that current experimental devices can not detect them.

References

- [1] M. R. Douglas and N. A. Nekrasov, Noncommutative Field Theory, *Rev. Mod. Phys.* **73** (2001) 977-1029
- [2] R. J. Szabo, Quantum Field Theory on Noncommutative Spaces, *Phys. Rept.* **378** (2003) 207-299
- [3] N. Seiberg and E. Witten, String Theory and Noncommutative Geometry, *JHEP* **9909** (1999) 032
- [4] A. Connes and M. Marcolli, A Walk in the Noncommutative Garden, arXiv:math.QA/0601054

- A. Connes, A Short Survey of Noncommutative Geometry, *J. Math. Phys.* **41** (2000) 3832-3866
- [5] A. Konechny and A. Schwarz, Introduction to M(atrix) Theory and Noncommutative Geometry, *Phys. Rept.* **360** (2002) 353-465;
 - [6] M. Chaichian *et al*, Noncommutative Standard Model: Model Building, *Eur. Phys. J. C* **29** (2003) 413-432
 - [7] A. Micu and M.M. Sheikh-Jabbari, Noncommutative Φ^4 Theory at Two Loops, *JHEP* **0101** (2001) 025
 - [8] S. Benczik et al, Short Distance vs. Long Distance Physics: The Classical Limit of the Minimal Length Uncertainty Relation, *Phys. Rev. D* **66** (2002) 026003
 - [9] B. Mirza and M. Dehghani, Noncommutative Geometry and the Classical Orbits of Particles in a Central force Potential, *Commun. Theor. Phys.* **42** (2004) 183-184
 - [10] K. Konishi, G. Paffuti, and P. Provero, Minimum physical length and the Generalized Uncertainty Principle in String Theory, *Phys. Lett. B* **234** (1990) 276
 - [11] A. Kempf et al, Hilbert space representation of the minimal length uncertainty relation, *Phys. Rev. D* **52** (1995) 1108
 - [12] M. S. El Naschie, The concepts of E-infinity: An elementary introduction to the Cantorian-fractal theory of quantum physics, *Chaos, Solitons and Fractals* **22** (2004) 495-511
 - [13] M. S. El Naschie, Silver Mean Hausdorff Dimension and Cantor Sets, *Chaos, Solitons and Fractals*, **4** (1994) 1861-1870
 - [14] S. M. Carroll, *An Introduction to General Relativity: Spacetime and Geometry*, Addison Wesley, 2004
 - [15] S. Benczik et al, Classical implication of the minimal length uncertainty relation , arXiv:math.QA/0209119




Figure 1: The effect of space noncommutativity on the stability of circular orbits of particles in a central force of the type $F(r) = -\frac{k}{r^n}$. Vertical axis shows the left hand side of relation (29). Note that in E -infinity theory, based on El Naschie bijection formula one can write $D_3 = \left(\frac{1}{d_c^{(0)}}\right)^{3-1} = 2 + \phi = \left(\frac{1}{\phi}\right)^2 = 2.618033969$ which is Hausdorff dimension of Penrose universe when time dimension is added. On the other hand this value is the intersection point of curve corresponding to noncommutative case with horizontal axis. This can be interpreted as $\left(\frac{1}{\phi}\right)^2 = 1 + 1.618033969$ where 1.618033969 is the golden mean[12,13].




Figure 2: The effects of space noncommutativity on the stability of circular orbits of particles in Yukawa potential. a) Commutative space, b) Noncommutative space with small L , and c) Noncommutative space with large L . Vertical axis shows the left hand side of relation (35). In the language of E -infinity theory, curves (a) and (b) intersect horizontal axis in 1.618033969 which is golden mean while curve (c) intersects horizontal axis at silver mean, 1.414213562 [12,13].




Figure 3: Commutative effective potential versus radius for different values of angular momentum(in arbitrary units).




Figure 4: The difference between commutative and noncommutative effective potential for $L = 1$ (in arbitrary units). Lower curve shows noncommutative case. For small values of L , the difference is not considerable.




Figure 5: The difference between commutative and noncommutative effective potential for $L = 5$. Lower curve shows noncommutative case. For large values of L , the difference is considerable.




Figure 6: The condition for stability of circular orbits for commutative and noncommutative Schwarzschild spaces. In commutative case, particles could have stable circular orbits for $r \geq 6GM$ while for noncommutative case $r \geq (6.139234690)GM$. Space noncommutativity increases the radius of stable circular orbits.